

- [0075] 17. Zemanick, E. T., Wagner, B. D., Robertson, C. E., Stevens, M. J., Szeffler, S. J., Accurso, F. J. et al. Assessment of airway microbiota and inflammation in cystic fibrosis using multiple sampling methods. *Ann Am Thorac Soc.* 2015; 12: 221-229.
- [0076] 18. Lee T J, Fu C H, Wang C H, et al. Impact of chronic rhinosinusitis on severe asthma patients. *PLoS ONE.* 2017; 12(2):e0171047.
- [0077] 19. Dixon A E, Kaminsky D A, Holbrook J T, Wise R A, Shade D M, Irvin C G. Allergic rhinitis and sinusitis in asthma: differential effects on symptoms and pulmonary function. *Chest.* 2006; 130(2):429-435.
- [0078] 20. Ramakrishnan V R, Hauser L J, Feazel L M, Ir D, Robertson C E, Frank D N. Sinus microbiota varies among chronic rhinosinusitis phenotypes and predicts surgical outcome. *J Allergy Clin Immunol.* 2015; 136(2):334-342. e1.
- [0079] 21. Mandavina M, Engen P A, LoSavio P S, et al. The nasal microbiome in patients with chronic rhinosinusitis: Analyzing the effects of atopy and bacterial functional pathways in 111 patients. *J Allergy Clin Immunol.* 2018; 142:287-290.
- [0080] 22. Fokkens W J, Lund V J, Mullol J, et al. EPOS 2012: European position paper on rhinosinusitis and nasal polyps 2012. A summary for otorhinolaryngologists. *Rhinology.* 2012; 50(1):1-12.
- [0081] 23. Cardenas P A, Cooper P J, Cox M J, et al. Upper airways microbiota in antibiotic-naïve wheezing and healthy infants from the tropics of rural Ecuador. *PLoS ONE.* 2012; 7(10):e46803.
- [0082] 24. Teo S M, Mok D, Pham K, et al. The infant nasopharyngeal microbiome impacts severity of lower respiratory infection and risk of asthma development. *Cell Host Microbe.* 2015; 17(5):704-715.
- [0083] 25. Huang Y J, Marsland B J, Bunyavanich S, et al. The microbiome in allergic disease: current understanding and future opportunities—2017 PRACTALL document of the American academy of allergy, asthma & immunology and the European academy of allergy and clinical immunology. *J Allergy Clin Immunol.* 2017; 139(4):1099-1110.
- [0084] 26. Navarro J, Rainisio M, Harms H K, et al. Factors associated with poor pulmonary function: cross-sectional analysis of data from the ERF. *European Epidemiologic Registry of Cystic Fibrosis.* *Eur Respir J.* 2001; 18(2):298-305.
- [0085] 27. David J, Bell R E, Clark G C. Mechanisms of disease: host-pathogen interactions between *Burkholderia* species and lung epithelial cells. *Front Cell Infect Microbiol.* 2015; 5:80.
- [0086] 28. Aanaes K. Bacterial sinusitis can be a focus for initial lung colonisation and chronic lung infection in patients with cystic fibrosis. *J Cyst Fibros.* 2013; 12(Suppl 2):51-520.
- [0087] 29. Lucas S K, Yang R, Dunitz J M, Boyer H C, Hunter R C. 16S rRNA gene sequencing reveals site-specific signatures of the upper and lower airways of cystic fibrosis patients. *J Cyst Fibros.* 2018; 17(2):204-212.

1. A method for treating sinusitis in a subject, the method comprising:

administering to the subject an effective amount of a composition that stimulates growth or activity of a genus of bacteria that is decreased relative to a control subject not having sinusitis and/or inhibiting growth or activity of a genus of bacteria that is increased relative to the control subject.

2. The method according to claim 1, wherein the composition comprises a bacterial composition.

3. The method according to claim 1, wherein the sinusitis is chronic rhinosinusitis (CRS).

4. The method according to claim 1, wherein the subject has asthma.

5. The method according to claim 1, wherein the composition stimulates growth or activity of the genus *Corynebacterium*.

6. The method according to claim 1, wherein the composition stimulates growth or activity of the genus *Peptoniphilus*.

7. The method according to claim 1, wherein the composition stimulates growth or activity of the genus *Corynebacterium* and the genus *Peptoniphilus*.

8. The method according to claim 1, wherein the composition comprises bacteria from the genus *Corynebacterium* and/or the genus *Peptoniphilus*.

9. The method according to claim 4, further comprising inhibiting growth or activity of the genus *Streptococcus*.

10. The method according to claim 4, further comprising inhibiting growth or activity of the genus *Burkholderia*.

11. The method according to claim 4, further comprising inhibiting growth or activity of the genus *Streptococcus* and the genus *Burkholderia*.

12. The method according to claim 1, wherein the composition comprises a bacterial composition that is harvested from a normal subject.

13. The method according to claim 12, wherein the bacterial composition is processed to isolate bacteria from the genus *Corynebacterium* and/or the genus *Peptoniphilus*.

14. The method according to claim 1, wherein the composition is administered intranasally.

15. The method according to claim 1, wherein the composition comprises a liquid, foam, cream, spray, powder, gel, or absorbent material.

16. The method according to claim 1, wherein a microbiome of the subject is analysed before administering the composition.

17. The method according to claim 1, wherein a microbiome of the subject is analysed after the administration of the composition.

18. A composition comprising a bacterial composition comprising bacteria from the genus *Corynebacterium* and/or the genus *Peptoniphilus*.

19. The composition according to claim 18, wherein the bacterial composition consists of bacteria from the genus *Corynebacterium* and the genus *Peptoniphilus*.

20. The composition according to claim 18, wherein the composition further comprises a second effective compound comprising an antibiotic.

21. The composition according to claim 18, wherein the composition further comprises an agent that is bacteriostatic or bactericidal to the genus *Streptococcus* and/or the genus *Burkholderia*.

\* \* \* \* \*